



JISAMAR®

Fertilizer with amino acids NPK 5-2-3 and Boron

GUARANTEED RICHNESS

Free amino acids	5,30% w/w
Total nitrogen (N)	5,80% w/w
Ammoniacal nitrogen (N)	0,40% w/w
Organic nitrogen (N)	1,10% w/w
Ureal nitrogen (N)	4,30% w/w
Phosphorus pentoxide (P ₂ O ₅) water soluble	2,50% w/w
Potassium oxide (K ₂ O) water soluble	3,80% w/w
Boron (B) water soluble	0,14% w/w



PROPERTIES

JISAMAR is a specially studied composition for foliar applications, containing free amino acids, such as proline, glycine and lysine, which have a direct effect on flowering and on the growth of the fruit.

JISAMAR also contains a 25,2% w/v of pure SEAWEED EXTRACT, *Ascophyllum Nodosum*, rich in phytohormones from vegetal origin, such as cytokinin, auxins, and gibberellins, which in combination with the supplied phosphorus are specially suitable for flowering, growth and ripening of the fruit.

DOSAGE AND INSTRUCTIONS FOR USE

JISAMAR is suitable for all kind of crops; especially where fruit quality and quality are important.

CROP	PERIOD OF APPLICATION	FOLIAR	ROOT
Vegetables	Post-transplantation and applications every 15-20 days	100-200 cc/hl	3-4 l/ha
Fruit trees	8-10 days before flowering, petal shedding and half fattening	150-250 cc/hl	4-5 l/ha
Citrus trees	Pre-flowering and petal shedding	100-150 cc/hl	3-4 l/ha
Olive trees	Sprouting, lump, fattening	150-250 cc/hl	4-5 l/ha
Grape trees	Pre-flowering and petal shedding	150-200 cc/hl	3-4 l/ha
Strawberry	10 to 20 days after rooting, with the appearance of first flowers and then with the phytosanitary treatments	150-200 cc/hl	3-4 l/ha
Banana	Applied with the phytosanitary treatments	100-200 cc/hl	3-4 l/ha

COMPATIBILITY

JISAMAR is compatible with all phytosanitary and nutritional products except those with very acid pH.

PRESENTATION

1; 5; 20; 220 & 1.000 L.